ECON42720 Causal Inference and Policy Evaluation 3 Potential Outcomes and Randomised Experiments

Ben Elsner (UCD)

About this Lecture

This lecture introduces the **potential outcomes framework**

- ▶ This framework allows us to think about causal effects in a structured way
- We learn why counterfactuals are important but also difficult to observe/construct
- ▶ We learn what biases can occur when we do not observe all confounders

One solution for causal inference: randomised experiments

- Under what conditions do they allow for the identification of a causal effect?
- How can a hypothetical experiment serve as a benchmark for other methods?

Lingo and Notation

The lingo of causal inference is borrowed from medical trials

- treatment is the intervention/variable whose effect we are interested in
- **treatment group** is the group of units that receives the treatment
- control group is the group of units that does not receive the treatment or receives a placebo
- outcome is the variable that is potentially affected by the treatment

We are after the causal effect: treatment $(D) \rightarrow$ outcome (Y)



This lecture is based on

Cunningham (2020), Chapter 4
Angrist & Pischke (2009), Chapter 2

Find more about the course on the course page

A common challenge in applied econometrics is to **separate a causal effect** from the **influence of third factors**



We often have a good (theoretical) idea why $D \rightarrow Y$

but X is a confounding factor

often the problem: we also have an idea what X could be...

but cannot observe it (notation: u for "unobservable'')



Common challenge: selection into treatment

In microeconomics we learn

- **people** make **rational choices**...
- ...as do firms
- ... as do governments

Problem: we do **not observe all the determinants** of these choices (i.e. u)

Examples for selection into treatment:

Going to the gym makes you healthier

- good reason to believe so
- but people who go to the gym are different from those who don't
- \blacktriangleright observed correlation \neq causation

Exporting boosts firm profitability

- good reason to believe so
- but exporters are different in many ways from non-exporters
- **b** observed correlation \neq causation

Problem: it is not easy to account for all confounding factors

- because we know what they are but can't observe them
- ▶ or because we don't know what they are (e.g. "common shocks' ')

The burden of proof is on the researcher

- If you make a causal statement, you need to make a convincing case
 but causality cannot be proven without assumptions...
- ... and these assumptions need to be believed

The Potential Outcomes Framework

We will now focus on binary treatments

Some notation from experimental studies

i is an index for the units in the population under study.

► *D_i* is the **treatment status**:

- ▶ $D_i = 1$ if unit *i* has been exposed to treatment,
- ▶ $D_i = 0$ if unit *i* has not been exposed to treatment.

• $Y_i(D_i)$ indicates the **potential outcome according to treatment**:

- $Y_i(1) \equiv Y_{i1}$ is the outcome in case of treatment,
- $Y_i(0) \equiv Y_{i0}$ is the outcome in case of no treatment.

The Potential Outcomes Framework (Rubin, 1974)

The observed outcome is then

$$Y_i = \begin{cases} Y_{i1} & \text{if } D_i = 1\\ Y_{i0} & \text{if } D_i = 0 \end{cases}$$

Switching equation

$$= D_i Y_{i1} + (1 - D_i) Y_{i0}$$

$$= Y_{i0} + \underbrace{(Y_{i1} - Y_{i0})}_{\text{treatment effect}} D_i$$

We are interested in the (individual) **treatment effect** $\Delta_i = Y_{i1} - Y_{i0}$.

The Fundamental Problem of Causal Inference

We cannot identify the individual treatment effect $\Delta_i = Y_{i1} - Y_{i0}$

This is logically impossible

- we either observe that a unit was treated or not
- but never both treatment statuses at the same time

Wanted: The Counterfactual

A critical ingredient to establish causality: the counterfactual

What would have happened to a unit if

- the treatment status $D \in \{0, 1\}$ was different?
- or, if D is continuous, the treatment intensity was different?

For units that were treated, we want to know $Y_{i0} \mid D_i = 1$. For units that were untreated, we want to know $Y_{i1} \mid D_i = 0$.

No counterfactual, no causal claim!

Finding a Counterfactual: The Average Treatment Effect (ATE)

Problem: we cannot observe the counterfactual

> at the individual level, the counterfactual is entirely hypothetical

Statistical solution: Average Treatment Effect (ATE) for a random unit $E(\Delta_i) = E(Y_{i1} - Y_{i0}) = E(Y_{i1}) - E(Y_{i0})$

If a random unit was treated, the expected difference in their outcome is the ATE

The Average Treatment Effect on the Treated (**ATT**)

The Average Treatment Effect on the Treated (ATT)

$$ATT = E(\Delta_i | D_i = 1) = E(Y_{i1} - Y_{i0} | D_i = 1) = E(Y_{i1} | D_i = 1) - E(Y_{i0} | D_i = 1).$$

Interpretation: Average difference in potential outcomes for those who were treated

The ATT is **useful for policy evaluation**: take units who were treated and think what their outcomes would be had they not been treated

The Average Treatment Effect on the Untreated (ATU)

Similarly, we can define the Average Treatment Effect on the Untreated (ATU) as:

$$ATU = E(\Delta_i | D_i = 0) = E(Y_{i1} - Y_{i0} | D_i = 0) = E(Y_{i1} | D_i = 0) - E(Y_{i0} | D_i = 0).$$

Interpretation: the average difference in potential outcomes for those who were not treated

The ATU can (sometimes) be **useful for policy evaluation**: take units who were untreated and think what their outcomes would be had they been treated

Treatment Effects: Numerical Example

Person	D	PO Treated Y ¹	PO Untreated Y ⁰	Causal Effect
1	1	80	60	20
2	1	75	70	5
3	0	70	60	10
4	1	85	80	5
5	0	75	70	5
6	0	80	80	0
7	0	90	100	-10
8	1	85	80	5

Here we observe the **potential outcomes** for each person. Some people are treated (D = 1), some are not (D = 0).

The Average Treatment Effect

Person	D	PO Treated Y ¹	PO Untreated Y ⁰	Causal Effect
1	1	80	60	20
2	1	75	70	5
3	0	70	60	10
4	1	85	80	5
5	0	75	70	5
6	0	80	80	0
7	0	90	100	-10
8	1	85	80	5

The ATE is the average of the last column: ATE = 5

The Average Treatment Effect on the Treated (ATT)

Person	D	PO Treated Y ¹	PO Untreated Y ⁰	Causal Effect
1	1	80	60	20
2	1	75	70	5
3	0	70	60	10
4	1	85	80	5
5	0	75	70	5
6	0	80	80	0
7	0	90	100	-10
8	1	85	80	5

The ATT is the average causal effect of those who were treated (green cells): ATT = 8.75

Person	D	PO Treated Y^1	PO Untreated Y ⁰	Causal Effect
1	1	80	60	20
2	1	75	70	5
3	0	70	60	10
4	1	85	80	5
5	0	75	70	5
6	0	80	80	0
7	0	90	100	-10
8	1	85	80	5

The Average Treatment Effect on the Untreated (ATU)

The ATU is the average causal effect of the units who were not treated (green cells): ATU = 1.25

Here we have ATT > ATE > ATU, which may indicate **selection into treatment**: those who benefit most from the treatment are most likely to take it

ATE/ATT/ATU: Which Parameter is most Relevant?

There is no clear answer to this question

ATE is the most general parameter

what if we give the average person/firm/unit a treatment

This is interesting for medical trials and for many policy questions

ATT is often interesting for policy evaluation

- take those who took up the policy
- what would have happened to them if they had not taken up the policy

ATU is sometimes interesting for policy evaluation

- ▶ We may be concerned about the people who did not take up the policy
- How would they be affected if they took up the policy

But: We need all three to interpret typical estimation results

Comparison by Treatment Status

In many applications, we want to estimate the ATE or the ATT

But we only observe

- Whether a unit was treated or not
- The actual outcome of the unit

Solution (?): **comparison of means/simple difference in outcomes**, which can be estimated from two samples of data (treatment and control group)

$$SDO = E(Y_i | D_i = 1) - E(Y_i | D_i = 0)$$

=
$$\underbrace{\frac{1}{N_T} \sum_i (y_i \mid d_i = 1)}_{\text{Treated units}} - \underbrace{\frac{1}{N_C} \sum_i (y_i \mid d_i = 0)}_{\text{"Control" units (untreated}}$$

Simple Difference in Outcomes (SDO)

Person	D	PO Treated Y ¹	PO Untreated Y ⁰	Causal Effect	
1	1	80	60	20	
2	1	75	70	5	
3	0	70	60	10	
4	1	85	80	5	
5	0	75	70	5	
6	0	80	80	0	
7	0	90	100	-10	
8	1	85	80	5	

We compare here the **observed outcomes** of the treated to the **observed outcomes** of the **untreated**

SDO = 81.25 - 77.5 = 3.75

Comparison by Treatment Status

Problem with SDO: treated and untreated units are not comparable

$$E(Y_i|D_i = 1) - E(Y_i|D_i = 0) = E(Y_{i1}|D_i = 1) - E(Y_{i0}|D_i = 0) = \underbrace{E(Y_{i1}|D_i = 1) - E(Y_{i0}|D_i = 1)}_{ATT} + \underbrace{E(Y_{i0}|D_i = 1) - E(Y_{i0}|D_i = 0)}_{\text{Selection bias}}$$

Selection bias: the potential outcomes would differ even if both groups were untreated

Note how we add and subtract $E(Y_{i0}|D_i = 1)$ here

Now Suppose We Want to Estimate the ATE

Note that we can write the ATE as: $ATE = \pi ATT + (1 - \pi)ATU$

The simple difference in outcomes yields:

$$E(Y_i|D_i = 1) - E(Y_i|D_i = 0) = ATE + \underbrace{E(Y_{i0}|D_i = 1) - E(Y_{i0}|D_i = 0)}_{\text{Selection bias}} + \underbrace{(1 - \pi)[ATT - ATU]}_{\text{HTE Bias}}$$

Our estimate of the ATE is contaminated by two biases:

selection bias

heterogeneous treatment effects (HTE) bias

Person	D	PO Treated Y^1	PO Untreated Y ⁰	Causal Effect
1	1	80	60	20
2	1	75	70	5
3	0	70	60	10
4	1	85	80	5
5	0	75	70	5
6	0	80	80	0
7	0	90	100	-10
8	1	85	80	5

Selection Bias: Genuine Difference between Treated and Untreated Units

The selection bias is the difference between the green and red cells.

In this case, $E(Y_{i0}|D_i = 1) - E(Y_{i0}|D_i = 0) = 72.5 - 77.5 = -5$

So in absence of the treatment, the untreated have more favorable outcomes than the treated $% \left({{{\mathbf{x}}_{i}}} \right)$

Selection Bias



Prince Charles

Male Born in 1948 Raised in the UK Married Twice Lives in a castle Wealthy and Famous



Ozzy Osbourne

Male Born in 1948 Raised in the UK Married Twice Lives in a castle Wealthy and Famous Selection Bias: Genuine Difference between Treated and Untreated Units

The selection bias is $E(Y_{i0}|D_i = 1) - E(Y_{i0}|D_i = 0)$

This means that, even without the treatment, the two groups would have different outcomes

Selection bias is very common! Whenever there is a **confounder**, we have selection bias

Heterogeneous Treatment Effects

HTE bias = $(1 - \pi) [ATT - ATU]$

Treatment effect may vary across units

People who take the treatment may be different from those who do not
 People with stronger treatment effects may be more likely to take the treatment

Heterogeneous treatment effects bias can even occur in a randomised experiment

- We can hardly ever enforce that people take the treatment
- Unlike in physics or chemistry, in the social sciences we can only encourage people to take the treatment
- We speak here of imperfect compliance

Heterogeneous Treatment Effects Bias



Source: Dall-E, OpenAl

Heterogeneous Treatment Effects Bias

Person	D	PO Treated Y ¹	PO Untreated Y ⁰	Causal Effect
1	1	80	60	20
2	1	75	70	5
3	0	70	60	10
4	1	85	80	5
5	0	75	70	5
6	0	80	80	0
7	0	90	100	-10
8	1	85	80	5

The **heterogeneous treatment effects bias** is the difference between the green and red cells weighted by the **probability of treatment**

HTE bias = $(1 - \pi) [ATT - ATU] = 0.5 \times [8.75 - 1.25] = 3.75$

SDO vs. ATT

In our example, SDO = 3.75

$$SDO = ATT + Selection Bias$$

= $8.75 - 5 = 3.75$

In our case, the **SDO under-estimates the ATT** by 5 because of selection bias. Without the treatment, treated units have less favourable outcomes than untreated units.

SDO vs ATE

$$SDO = ATE +$$
 Selection Bias + HTE Bias
= 5 - 5 + 3.75
= 3.75

In our case, the SDO also under-estimates the ATE. This is the result of two biases going in opposite directions. The overall bias is negative.

The previous slides have shown that

- a simple (a.k.a. naive) comparison of treated and untreated units yields a biased estimate of the ATE and ATT
- It is not random whether a unit is treated or not or chooses to get treated or not
- This choice leads to selection bias and heterogeneous treatment effects bias
- A randomised experiment can eliminate selection bias but not necessarily the HTE bias

So when Is the SDO Unbiased?

The SDO is unbiased (SDO = ATE) under the **Independence Assumption**

$$(Y_1, Y_0) \perp D \tag{1}$$

The potential outcomes are independent of the treatment

- Treated and untreated individuals have the same average potential outcomes E(Y₀) and E(Y₁)
- I.e. they are statistically similar
- And no one selects into treatment because of their treatment effect

So when Is the SDO Unbiased?

We can show that the selection bias and HTE bias are zero under the independence assumption

The independence assumption implies

$$E[Y_1 | D = 1] - E[Y_1 | D = 0] = 0$$

$$E[Y_0 | D = 1] - E[Y_0 | D = 0]$$

Selection bias

Independence also implies that ATU = ATT

$$ATT - ATU = E[Y_1 | D = 1] - E[Y_0 | D = 1]$$
$$- E[Y_1 | D = 0] + E[Y_0 | D = 0]$$
$$= 0$$

Implication for research practice:

We need to understand the sources of bias in our context

We cannot test directly for the presence of bias

no matter how much big data enthusiasts make us believe that we can

We need to be humble about **what parameters we can identify with our data** - Example: ATT does not suffer from HTE bias, but is it relevant?

Comparison by Treatment Status

In most cases, we run regressions rather than comparing means

Take expectations conditional on D_i :

$$E(Y_i|D_i = 1) = \alpha + \beta + E(\varepsilon_i|D_i = 1)$$

$$E(Y_i|D_i = 0) = \alpha + E(\varepsilon_i|D_i = 0)$$

Comparison by Treatment Status

Your regression will estimate a coefficient $\hat{\beta}$ that is a combination of a treatment effect (ATE or ATT) plus a bias

The OLS estimate of β is

$$E(Y_i|D_i = 1) - E(Y_i|D_i = 0) = \underbrace{\beta}_{\text{Treatment Effect}} + \underbrace{E(\varepsilon_i|D_i = 1) - E(\varepsilon_i|D_i = 0)}_{\text{bias}}.$$

The RHS is either

ATT + Selection Bias
 ATE + Selection Bias + HTE Bias
 Both are equivalent

We should always assume that the bias $\neq 0$

What if We Control for X

Person	D	Female (X)	PO Treated Y ¹	PO Untreated Y ⁰	Causal Effect
1	1	1	80	60	20
2	1	1	75	70	5
3	0	1	70	60	10
4	1	0	85	80	5
5	0	1	75	70	5
6	0	0	80	80	0
7	0	0	90	100	-10
8	1	0	85	80	5

Controlling for X

If X is binary, we compare the outcomes conditional on X

This means that

- We compare the outcomes of treated and untreated units within each group defined by X
- Our estimator is the weighted average of these differences

Consider p = Pr(x = 1) and q = Pr(x = 0), with p + q = 1

$$\begin{split} E(Y_i|D_i = 1, X) &- E(Y_i|D_i = 0, X) \\ &= p \left[E(Y_{i1} \mid D_i = 1, x = 1) - E(Y_{i1} \mid D_i = 0, x = 1) \right] \\ &+ q \left[E(Y_{i1} \mid D_i = 1, x = 0) - E(Y_{i1} \mid D_i = 0, x = 0) \right] \\ &= p \left[ATT_{x=1} + E(Y_{i0} \mid D_i = 1, x = 1) - E(Y_{i0} \mid D_i = 0, x = 1) \right] \\ &+ q \left[ATT_{x=0} + E(Y_{i0} \mid D_i = 1, x = 0) - E(Y_{i0} \mid D_i = 0, x = 0) \right] \end{split}$$

What if We Control for X

Suppose \wedge is binary, for example male and remain	Suppose	Χ	is	binary,	for	example	male	and	female
---	---------	---	----	---------	-----	---------	------	-----	--------

Person	D	Female (X)	PO Treated Y ¹	PO Untreated Y ⁰	Causal Effect
1	1	1	80	60	20
2	1	1	75	70	5
3	0	1	70	60	10
4	1	0	85	80	5
5	0	1	75	70	5
6	0	0	80	80	0
7	0	0	90	100	-10
8	1	0	85	80	5

►
$$SDO_{X=1} = 77.5 - 65 = 12.5$$

► $SDO_{X=0} = 85 - 90 = -5$
► $SDO = \frac{1}{2} \times 12.5 + \frac{1}{2} \times (-5) = 3.75$

Controlling for X

In our example, E(D | X) = E(D), i.e. *D* is independent of *X*

▶ This means that controlling for *X* does not affect the estimate of the SDO

But often, $E(D | X) \neq E(D)$ because X is a confounder

That's exactly when we want to control for X
If X is the only confounder, controlling for X eliminates the selection bias

We can identify the ATE under the Conditional Independence Assumption

 $(Y_1, Y_0) \perp D \mid X$

Experiments provide a clean way to eliminate selection bias

Even if in most cases it is **not possible to run experiments**, a **clean experiment** serves as the **benchmark** for all the methods in this course

Idea: the closer we get to the **ideal experimental setting**, the closer we get to estimating a **causal effect**

Basic idea: randomly assign treatment across units

Two conditions have to be fulfilled:

assignment to treatment and control group is random
 there is full compliance with the treatment (plus: no attrition)

In that case, the treatment and control group are statistically identical

Formally (T: treatment group, C: control group)

$$E\{Y_{i0}|i \in C\} = E\{Y_{i0}|i \in T\}$$

 and

$$E\{Y_{i1}|i \in C\} = E\{Y_{i1}|i \in T\}.$$

both groups have the same potential outcomes in expectation

Therefore, we can obtain an unbiased and consistent estimate of the ATE

$$E(\Delta_i) = E(Y_{i1} - Y_{i0}) = E\{Y_{i1} | i \in T\} - E\{Y_{i0} | i \in C\}.$$

 \Rightarrow comparison of means is meaningful (causal effect in expectation)

 \Rightarrow Randomisation solves the fundamental problem of causal inference

But what if not everyone complies with the treatment?

Examples

Not every job seeker who is offered training takes part
 Not every person eligible for a medical card/social benefits/etc applies

$$E\{Y_{i1}|i \in T\} - E\{Y_{i0}|i \in C\} \neq ATE$$

As we will learn later in the course, all is not lost

The resulting treatment effect can still be meaningful
 ... as long as treatment is random

An important assumption underlying causal inference in experiments

SUTVA: Stable Unit Treatment Value Assumption

In plain English

treatment of one unit must not affect the potential outcomes of another

▶ i.e. no spillovers, general equilibrium effects, etc

SUTVA is an untestable assumption

Examples for violations:

- information leaks from treatment to control group
- ► large-scale job training programs ⇒ GE effects
- health interventions (capacity constraints in medical facilities)

Implications:

- One reason why small interventions don't scale up
- Important to choose the right control group

Cookbook for Analyzing a Randomised Experiment

- 1) Explain experimental design in detail, in particular
 - How was the randomisation carried out
 - Discuss compliance (or likely non-compliance)
 - Argue why SUTVA holds

2) Show balancing tests based on pre-treatment characteristics

- Do treatment and control group differ before the experiment?
- Use pre-treatment outcomes if possible
- Use other pre-treatment characteristics that may predict selection into treatment
- If you use regression analysis in the paper, regress the pre-treatment x on the treatment
- Never use post-treatment outcomes (NEVER EVER!)

$$x_i = \delta_1 + \delta_2 D_i + \eta_i$$

Cookbook for Analyzing a Randomised Experiment

3) Show and discuss results

Compare means across treatment and control groups

Add **pre-treatment characteristics** X_i as controls

 $y_i = \alpha + \beta D_i + \boldsymbol{X_i \gamma} + u_i$

Why add controls?

Additional test if randomisation worked (β should not change)
 Estimates become more precise (less noise in the model ⇒ lower SE)

Research Question: does class size matter to student learning? (Krueger, 1999)

Implemented on cohort of kindergartners (i.e. senior infants) in 1985/86 in Tennessee.

Lasted 4 years, then everyone went back to regular size class.

Three treatments

- 1. Small class 13-17
- 2. Regular class 22-25
- 3. Regular class with Aide

Schools had to have at least 3 classes to participate.

Entering cohort randomly assigned to class type. Teachers also randomly assigned.

I.e. randomisation was carried out within schools

Test scores measured towards end of each school year in March.

Balancing tests: do treatment and control groups have the **same pre-treatment** characteristics?

		Class	P-value for equalit	
Variable	Small	Regular	Regular/Aide	across groups
Free lunch	.47	.48	.50	.09
White/Asian	.68	.67	.66	.26
Age in 1985	5.44	5.43	5.42	.32
Attrition rate	.49	.52	.53	.02
Class size in kindergarten	15.10	22.40	22.80	.00
Percentile score in kindergarten	54.70	48.90	50.00	.00

TABLE 2.2.1 Comparison of treatment and control characteristics in the Tennessee STAR experiment

Notes: Adapted from Krueger (1999), table I. The table show means of variables by treatment status for the sample of students who entered STAR in kindergarten. The *P*-value in the last column is for the *P*-test of equality of variable means across all three groups. The free lunch variable is the fraction receiving a free lunch. The percentile score is the average percentile score on three Stanford Achievement Tests. The attrition rate is the proportion lost to follow-up before completing third grade.

- They are similar in age, race and SES
- Attrition rates are similar
- Treatment and outcomes differ

Krueger (1999) uses a regression to analyse the experiment

The most comprehensive specification includes pre-treatment characteristics and school fixed effects

$$y_{is} = lpha + eta D_{is} + oldsymbol{X}'_{is} oldsymbol{\gamma} + \delta_s + u_{is}$$

Why include fixed effects?

- Randomisation was carried out within schools
- Children in different school types may react differently to the treatment
- ▶ The FE estimator compares children within the same school

Terrer de la colorada	TABLE	2.2.2	1	
Experimental estimate	s of the e	effect of c	class size on	test scores
Explanatory Variable	(1)	(2)	(3)	(4)
Small class	4.82 (2.19)	5.37 (1.26)	5.36 (1.21)	5.37 (1.19)
Regular/aide class	.12 (2.23)	.29 (1.13)	.53 (1.09)	.31 (1.07)
White/Asian			8.35 (1.35)	8.44 (1.36)
Girl			4.48 (.63)	4.39 (.63)
Free lunch		_	-13.15 (.77)	-13.07 (.77)
White teacher		—	_	57 (2.10)
Teacher experience				.26 (.10)
Teacher Master's degree				-0.51 (1.06)
School fixed effects	No	Yes	Yes	Yes
R ²	.01	.25	.31	.31

Notes: Adapted from Krueger (1999), table V. The dependent variable is the Stanford Achievement Test percentile score. Robust standard errors allowing for correlated residuals within classes are shown in parentheses. The sample size is 5,681.

Result: smaller classes are more effective

Experimental design appears valid

Balancing tables point to clean randomisation
 Estimates not affected by controls for pre-treatment characteristics

The same cookbook approach can be used for non-experimental studies

Discrete vs. continuous treatment

So far, we discussed experiments with a discrete treatment

The same assumptions apply to experiments with a continuous treatment

- The treatment intensity varies across units
- treatment intensity is randomly assigned

Possibilites for researchers:

Discretise and compare mean outcomes (e.g. above/below median)
 Estimate marginal effect in a regression

Randomised Experiments as Templates

Randomised experiments are often difficult or impossible to conduct

But they serve as a **template for estimating causal effects** in non-experimental settings

Any study that claims to estimate a causal effect should

- explain what the treatment is
- and under what conditions the assignment of the treatment (intensity) is as good as random
- ... even if the world is not perfect, there is no harm thinking about the perfect

My advice (after 100+ referee reports, several published papers and many rejections): ignore the experimental template at your own peril

References

Angrist, Joshua, & Pischke, Jörn-Steffen. 2009. Mostly Harmless Econometrics - An Empiricist's Companion. Princeton University Press. Cunningham, Scott. 2020. Causal Inference: The Mixtape. Yale University Press.

Krueger, Alan B. 1999. Experimental Estimates of Education Production Functions. *The Quarterly Journal of Economics*, **114**(2), 497–532. Rubin, Donald. 1974. Estimating Causal Effects of Treatments in Randomized and Nonrandomized Studies. *Journal of Educational Psychology*, **66**(5).

Appendix

Group Work I

Suppose you want to study the impact of medical cards on health outcomes in Ireland. Medical cards are means-tested and given to people with low income. Card holders can access medical care for free. You have access to administrative data covering all adults in Ireland, including information on whether they ever held a medical card and their health outcomes.

Tasks:

- 1. Explain: What is the treatment in this case?
- 2. Intuitively, explain what the ATE, ATU, ATT are. Which of these parameters is most relevant in a policy evaluation in this context?
- 3. What are potential sources of bias here?

Group Work II

Consider the independence assumption

$$(Y_1, Y_0) \perp D$$

Evaluate the following statements (i.e. are they implied by the independence assumption)?

1.
$$E[Y_1 | D = 1] - E[Y_0 | D = 0] = 0$$

2. $E[Y_1 | D = 1] - E[Y_0 | D = 1] = 0$
3. $E[Y_1 | D = 1] - E[Y_1 | D = 0] = 0$

Group Work III

Suppose you want to study the impact of local police presence on crime in Dublin.

Questions:

- What would be the ideal experiment to run here?
- Why may this experiment not be feasible?
- If you can't run an experiment, what else could you do to get at the causal effect?

Derivation of SDO Decomposition

$$E(Y^{1} | D = 1) - E(Y^{0} | D = 0)$$

= $E(Y^{1} | D = 1) - E(Y^{1} | D = 0) + E(Y^{1} | D = 0) - E(Y^{0} | D = 0)$
= $\underbrace{E(Y^{1} | D = 1) - E(Y^{1} | D = 0)}_{ATT} + \underbrace{E(Y^{1} | D = 0) - E(Y^{0} | D = 0)}_{Selection Bias}$

Derivation of SDO Decomposition

Now consider $ATE = \pi ATT + (1 - \pi)ATU$

$$egin{aligned} \mathsf{ATE} &= \pi \mathsf{ATT} + (1-\pi) \mathsf{ATU} \ &= \pi \mathsf{ATT} + \mathsf{ATT} - \mathsf{ATT} + (1-\pi) \mathsf{ATU} \end{aligned}$$

Re-arranging yields

$$ATT = ATE + (1 - \pi)ATT - (1 - \pi)ATU$$

= $ATE + (1 - \pi)[ATT - ATU]$

Derivation of SDO Decomposition

We can plug this into the SDO decomposition and get

$$E(Y^{1} | D = 1) - E(Y^{0} | D = 0)$$

$$= ATE$$

$$+ \underbrace{E(Y^{1} | D = 0) - E(Y^{0} | D = 0)}_{\text{Selection Bias}}$$

$$+ \underbrace{(1 - \pi)[ATT - ATU]}_{\text{HTE Bias}}$$



.

benjamin.elsner@ucd.ie



www.benjaminelsner.com



Sign up for office hours



YouTube Channel



@ben_elsner

LinkedIn

Contact

Prof. Benjamin Elsner

University College Dublin School of Economics Newman Building, Office G206 benjamin.elsner@ucd.ie

Office hours: book on Calendly